

Classical Stringy Black Holes Modify the Thermal Spectrum

B. Harms and Y. Leblanc

Department of Physics and Astronomy, The University of Alabama,

Tuscaloosa, AL 35487-0324

Non-local (α') corrections to Schwarzschild black holes are shown to invalidate the thermodynamical interpretation of black holes. In particular, the canonical and Bekenstein-Hawking temperatures are not equal. The particle number density of fields quantized in the (α' modified) black hole background is no longer thermal. In the non-perturbative region $\alpha' \rightarrow \infty$ (or $M \rightarrow 0$), an analytic continuation to the number density is shown to vanish exponentially.

PACS numbers: 04.60.+n, 11.17.+y, 97.60.Lf

In a previous paper [1], we calculated the stringy (α') classical corrections to the 4-dimensional Schwarzschild black hole metric and studied their effects on Hawking's black hole thermodynamics. It was found [1] that both the area law and the first law of black hole thermodynamics are invalidated by the α' -perturbation corrections. The breakdown of the first law meant the inequality of the canonical and the Bekenstein-Hawking temperature [2,3].

In this letter we make use of the above findings to provide a simple calculation of the quantum particle number density in an (α' -) modified black hole background. The so-called back-reaction (quantum gravity) effects are neglected here. We shall find that the breakdown of the first law implies that the particle spectrum is no longer thermal and that, on the basis of a reasonable speculation on the strong (non-perturbative) α' -limit ($\alpha' \rightarrow \infty$) or small black hole mass ($M \rightarrow 0$), the number density vanishes at least exponentially. This result is in agreement with the identification of extreme neutral black holes as massless elementary particles [4–7], contrary to the usual thermal interpretation (with infinite temperature).

The starting point of our exposition is the well known α' -modified Einstein field equations [8],

$$R_{\mu\nu} + \frac{\alpha'}{2} R_{\mu\kappa\lambda\tau} R_{\nu}^{\kappa\lambda\tau} + O(\alpha'^2) = 0 , \quad (1)$$

of a 26-dimensional bosonic string theory. These equations are obtained from the requirement of the vanishing of the tree-level dilaton tadpole in a curved background [9]. As before [1] we assume that all but four spacetime dimensions have been compactified with the internal space volume taken equal to unity for simplicity.

Applying perturbation theory about the small Regge slope α' ($\alpha' = \frac{1}{2\pi T}$ where T is the string tension), the unperturbed metric chosen to be the 4-dimensional Schwarzschild metric, Eq(1), yields the following solution for the modified black hole spacetime [1]

$$ds^2 = -\lambda^2(r)dt^2 + \lambda^{-2}(r)dr^2 + r^2(d\theta^2 + \sin^2\theta d\phi^2) , \quad (2)$$

where,

$$\lambda^2(r) = 1 - \frac{2M}{r} - \frac{2M^2\alpha'}{r} \left(\frac{1}{r_+^3} - \frac{1}{r^3} \right) + O(\alpha'^2) , \quad (3)$$

in which $r_+ = 2M$ is the black hole horizon.

An interesting result of our calculations is that, at least perturbatively, the non-local α' -effects do not alter the spacetime topology and the horizon remains unshifted.

According to Gibbons and Hawking [10], black hole thermodynamics can be studied by analytically continuing the Schwarzschild metric to the Euclidean domain, the period of the compact Euclidean time representing the inverse (Bekenstein - Hawking) temperature of black hole systems now endowed with thermal attributes. In particular, the Euclidean action evaluated at the black hole solution has received an interpretation in terms of thermodynamical entropy, a result justified by the following WKB expression for the black hole canonical partition function,

$$Z(\beta_H) \sim e^{-S_E/\hbar} . \quad (4)$$

In a series of recent papers however [4–7], we have shown the inconsistency of the thermal (canonical) ensemble for black holes and have pointed out the inequivalences between the canonical and microcanonical ensembles (as reflected by a negative microcanonical specific heat). In particular, the equilibrium state was shown to be very inhomogeneous, and thus very far from a thermal state [4–7].

Nevertheless, we wish here to play the devil's advocate and analyze the effects of stringy non-locality from the viewpoint of the thermodynamical interpretation.

In Euclidean spacetime the important requirement of the vanishing of the conical (deficit angle) singularity yields directly the period (inverse Hawking temperature) of the compact Euclidean time dimension [1],

$$\beta_H = 2\pi \left[\lambda(r) \frac{d}{dr} \lambda(r) \right]_{r=r_+}^{-1} \simeq 8\pi M \left(1 + \frac{3\alpha'}{8M^2} + O(\alpha'^2) \right) . \quad (5)$$

The entropy S_E was calculated in Ref. [1]. To order α'^2 we found,

$$S_E = \beta_H \left(\frac{M}{2} + \frac{\alpha'}{8M^2} + O(\alpha'^2) \right)$$

$$\begin{aligned}
&= 4\pi M^2 \left(1 + \frac{5\alpha'^2}{8M^2} + O(\alpha'^2)\right) \\
&= \frac{A}{4} \left(1 + \frac{5\alpha'}{8M^2} + O(\alpha'^2)\right) ,
\end{aligned} \tag{6}$$

in which we used Eq.(5) and where $A = 4\pi r_+^2$, i.e. the area of the horizon.

The canonical temperature of the gas is given by the usual thermodynamical relation [1]

$$\beta = \frac{dS_E}{dM} \simeq 8\pi M(1 + O(\alpha'^2)) . \tag{7}$$

Two important implications follow from these results. First Eq.(6) clearly shows a modification of the area law of black hole thermodynamics. Second, and perhaps more importantly, the Bekenstein-Hawking temperature, Eq.(5), is not equal to the above canonical temperature of the supposedly thermal equilibrium state of the gas,

$$\beta \neq \beta_H . \tag{8}$$

This is an explicit breakdown of the first law of black hole thermodynamics.

Canonical quantization of, say, a massless scalar field in the (α' - modified) classical black hole background (one neglects the back reaction here) should, pending accuracy of the thermodynamical description, yield a Planckian distribution for the particle number density at the equilibrium canonical temperature β . However, because the α' - effects do not shift the horizon or alter the topology (perturbatively), the scalar particle Green's functions turn out to be periodic (in the sense of the Kubo- Martin-Schwinger condition) with period β_H . These are temperature Green's functions and yield a Planckian distribution for the particle number density with temperature β_H^{-1} . Therefore,

$$\begin{aligned}
n(\omega) &= \frac{1}{e^{\beta_H \omega} - 1} \\
&\simeq \frac{1}{e^{\beta \omega} - 1} - \delta\beta \frac{\omega e^{\beta \omega}}{(e^{\beta \omega} - 1)^2} + O(\delta\beta^2) ,
\end{aligned} \tag{9}$$

where,

$$\delta\beta \equiv \beta_H - \beta \simeq \frac{3\pi\alpha'}{M} + O(\alpha'^2) . \tag{10}$$

Eq.(9) shows that the stringy effects modify the thermal distribution. Quantum gravity (back reaction) effects are widely believed to modify the thermal spectrum even more drastically. One can imagine loop effects effectively creating a cosmological constant term in the action. Such a term would completely change the topology of the black hole spacetime and consequently the particle number density will very likely not be Planckian. Black holes with a cosmological constant usually have two horizons and a metric described by the de Sitter-Schwarzschild solution.

Eq.(10) shows that small (light) black holes belong to the non-perturbative domain ($\alpha' \rightarrow \infty$). From the viewpoint of string theory, a duality transformation from strong to weak string tension may prove useful in the investigation of black holes. Here, however, we can only speculate on the behavior of the theory of strong α' . As noticed in our previous work [1] and as is apparent from Eq.(3), higher order α' - corrections actually worsen the $r \rightarrow 0$ (ultraviolet) behavior of the black hole spacetime. A conjecture called the Limiting Curvature Hypothesis (LCH) [11] however requires that the full gravity theory be singularity free. This would imply that, although each individual term in the α' - expansion is singular, the series should be finite. The above hypothesis is based on the seemingly sensible physical requirement of geodesic completeness.

An analytic continuation to strong α' ($\alpha' \rightarrow \infty$) or equivalently small black holes ($M \rightarrow 0$) was suggested in Ref. [1]. In this continuation Eq.(3) is the small α' -expansion of the expression

$$\lambda^2(r) = 1 - \frac{2M}{r} e^{\alpha' M (\frac{1}{r_+^3} - \frac{1}{r^3})} . \quad (11)$$

The above form effectively regularizes the black hole singularity at the origin as gravity is shown to vanish there (asymptotic freedom). An α' - modified graviton propagator can be derived from such a spacetime and was shown to yield a UV-finite quantum theory of gravity [1].

In the thermodynamical interpretation, it is interesting to speculate that Eq.(5) for the Hawking temperature could be analytically continued to the small black hole domain as

follows [1],

$$\beta_H = 8\pi M e^{3\alpha'/8M^2} . \quad (12)$$

Contrary to the usual case where α' - effects are neglected, we now find that extreme black holes should have zero Hawking temperature. Notice that Eq.(12) does not follow from the choice of Eq.(11) for the metric. Inserting Eq.(12) into the formula (Eq.(9)) for the particle number density, one would find the following asymptotic behavior,

$$n(\omega) \sim \exp(-8\pi M \omega e^{3\alpha'/8M^2}) \quad ; (M \rightarrow 0) , \quad (13)$$

a result to be compared with the following behavior for large black holes,

$$n(\omega) \sim \exp(-8\pi M \omega) \quad ; (M \rightarrow \infty) . \quad (14)$$

The reader will by now surely be aware that the continuation Eq.(11) of Eq.(3) is not unique. It maybe interesting to note that any generalization of Eq.(3) of the form

$$\lambda^2 = 1 - \frac{2M}{r} \sum_{n=0}^{\infty} C_n \alpha'^n M^n \left(\frac{1}{r_+^3} - \frac{1}{r^3} \right)^n , \quad (15)$$

with $C_0 = C_1 = 1$ as required by Eq.(3), leads to the same Hawking temperature,

$$T_H = \frac{1}{8\pi M} \left[1 - \frac{3\alpha'}{8M^2} \right] . \quad (16)$$

Thus the Hawking temperature goes to zero for $M \rightarrow \infty$ and $M^2 = 3\alpha'/8$. Below the latter value T_H approaches $-\infty$. Clearly, explicit calculation of the higher order corrections in α' are badly needed.

Finally, although the considerations of this letter clearly show that the exact thermal nature of the particle spectrum in a black hole background is spoiled by the non-local stringy effects, the loss of coherence problem is not resolved by the inclusion of non-local effects. The resolution of this problem must reside in the formulation of a self-consistent theory of quantum gravity.

ACKNOWLEDGMENTS

This research was supported in part by the U.S. Department of Energy under Grant No. DE-FG05-84ER40141.

REFERENCES

- [1] B. Harms and Y. Leblanc, University of Alabama preprint UAHEP-935.
- [2] J.D. Bekenstein, Phys. Rev. **D7**, 2333(1973); Phys. Rev. **D9**, 3292(1974).
- [3] S.W. Hawking, Comm. Math. Phys. **43**, 199(1975); Phys. Rev. **D13**, 191(1975).
- [4] B. Harms and Y. Leblanc, Phys. Rev. **D46**, 2334(1992).
- [5] B. Harms and Y. Leblanc, Phys. Rev. **D47**, 2438(1993).
- [6] B. Harms and Y. Leblanc, to appear in the *Proceedings of the Texas/PASCOS Conference*, University of California, Berkeley, CA (December 1992).
- [7] B. Harms and Y. Leblanc, to appear in the *Proceedings of SUSY-93*, Northeastern University, Boston, MA (March 1993).
- [8] M.B. Green, J.H. Schwarz and E. Witten, *Superstring Theory, Vol. I*, Cambridge University Press, Cambridge, 1987.
- [9] H. Nishimura, M. Tabuse and M. Takama, Phys. Lett. **B203**, 251(1988).
- [10] G.W. Gibbons and S.W. Hawking, Phys. Rev. **D15**, 2752(1977).
- [11] R. Brandenberger, V. Mukhanov and A. Sornborger, BROWN-HET-891 (1993), unpublished. And references therein.